

Fluorescence Turn-On Sensor for Cyanide Based on a Cobalt(II)–Coumarinylsalen Complex


Jae Han Lee,[†] A Reum Jeong,[†] Ik-Soo Shin,[†] Hae-Jo Kim,^{*,‡} and Jong-In Hong^{*,†}

Department of Chemistry, College of Natural Sciences, Seoul National University, Seoul 151-747, Korea, and Department of Chemistry, Kyonggi University, Suwon 443-760, Korea

jihong@snu.ac.kr; haejkim@kgu.ac.kr

Received December 10, 2009

ABSTRACT



A Co(II)–salen based fluorescent sensor (1-Co) that can selectively recognize cyanide anions in 1:2 binding stoichiometry over other anions has been developed. 1-Co displayed fluorescence enhancement upon the addition of cyanide owing to the interruption of photoinduced electron transfer from the coumarin fluorophore to the cobalt(II) ion. A general regression method was developed to calculate the binding constants in the 1:2 binding system, through which the 1:2 binding between 1-Co and cyanide anions was estimated to be in the range of micromolar dissociation constants.

Optical sensors for anions have gained considerable attention due to their applicability to the detection of toxic anions, monitoring of enzyme reactions related to anions, and bioimaging of anions.¹ Cyanide anions are known to inhibit the process of cellular respiration in mammals by interacting strongly with a heme unit in the active site of cytochrome

a₃.² The high level of intracellular calcium ion concentration is caused by the uptake of cyanide anions and triggers a cascade of enzymatic reactions to increase the level of reactive oxygen species, which finally inhibits the antioxidant defense system.³ The widespread use of cyanide anions in industrial settings (1.5 million tons per year) and potential threats for terrorism⁴ also increase the significance to sense cyanide anions by a simple and fast luminescence method. Most cyanide sensors operate on the basis of a 1:1 binding

[†] Seoul National University.

[‡] Kyonggi University.

(1) (a) Brzózka, Z. *Comprehensive Supramolecular Chemistry*; Atwood, J. L., Davies, J. E. D., MacNicol, D. D., Vögtle, F., Suslick, K. S., Eds.; Pergamon: Oxford, 1996; pp 187–212. (b) *Chemosensors of Ion and Molecular Recognition*; Desvergne, J.-P., Czarnik, A. W., Eds.; Kluwer: Dordrecht, 1997; Vol. 492. (c) Schmidtchen, F. P.; Berger, M. *Chem. Rev.* **1997**, *97*, 1609–1646. (d) Beer, P. D. *Acc. Chem. Res.* **1998**, *31*, 71–80. (e) *Supramolecular Chemistry for Anions*; Binachi, K., Bowman-James, K., Garcia-Espana, E., Eds.; New York, 1997. (f) Lehn, J.-M. *Supramolecular Chemistry, Concepts and Perspectives*; VCH: Weinheim, 1995.


(2) (a) Warburg, O. *Hoppe-Seyler's Z. Physiol. Chem.* **1911**, *76*, 331–346. (b) Kellin, D. *Proc. R. Soc. London, Ser. B* **1929**, *104*, 206–251. (c) Vennesland, B.; Comm, E. E.; Knowles, C. J.; Westly, J.; Wissing, F. *Cyanide in Biology*; Academic Press: London, 1981.

(3) (a) Johnson, J. D.; Meisenheimer, T. L.; Isom, G. E. *Toxicol. Appl. Pharmacol.* **1986**, *84*, 464–469. (b) Ardel, B. K.; Borowitz, J. L.; Isom, G. E. *Toxicology* **1989**, *56*, 146–154.

mode whether they are transition metal complexes⁵ or nonmetal probes.⁶ The 1:2 binding system, however, does closely resemble nature's binding or recognition systems like antibody–antigen⁷ and protein–metal ion binding.⁸ We thought that 1:2 binding between a metal complex and a cyanide anion would provide a useful sensing mechanism. Herein, we report a Co(II)–salen based fluorescent sensor (**1**·Co) bearing coumarin derivatives on the 1,2-positions of the ethylenediamine part of the salen ligand and cobalt(II) ion as a fluorescence quencher. The cobalt(II)–salen complex exhibited selective and tight binding toward cyanide anions on the basis of a 1:2 host–guest binding mode.

In addition, we developed a general method to determine the binding constants for a 1:2 binding system using a C program. Generally, a 1:2 binding system usually involves difficulties in quantifying binding properties such as binding constant and stoichiometry compared with a 1:1 binding system. We expected that the calculation method developed for the current 1:2 binding system will establish a general solution for quantification of the binding properties in multiple binding systems.

Compound **1** was prepared by the treatment of 2,2'-dihydroxyphenylene ethylenediamine with 2 equiv of 7-diethylaminocoumarin-3-carboxaldehyde through a diaza-Cope rearrangement reaction.⁹ The metal complex, **1**·Co, was obtained by mixing **1** with cobalt(II) acetate in the presence of triethylamine (TEA), followed by recrystallization of the complex in methanol and dichloromethane (Scheme 1).¹⁰



The effect of the metal coordination on the emission spectra was measured by fluorescence spectroscopy: the cobalt–salen complex (**1**·Co) showed weak fluorescence emission compared with ligand **1** because there seems to exist an intramolecular photoinduced electron transfer (PeT) from the coumarin moiety to the cobalt cation of **1**·Co. We expected that the addition of cyanide anions would cause an alteration in HOMO and LUMO energy levels and lead to a significant change in the fluorescence intensity of **1**·Co by blocking the PeT pathway (Figure 1).

As expected, we found that the fluorescence of the **1**·Co complex was significantly enhanced upon the addition of cyanide anions. 1:2 Complexation between **1**·Co and the cyanide anion was confirmed by high-resolution mass spectral analysis: observed m/z 803.2341, calculated 803.2341




Figure 1. Complexation of the cyanide ion with **1**·Co.

for [**1**·Co(CN[−])₂]. The effect of anions on the emission spectra of **1**·Co was examined by adding various anions to the cobalt–salen complex in acetonitrile. **1**·Co exhibited a significant fluorescence enhancement upon the addition of cyanide, whereas other anions showed no detectable fluorescence changes (Figure 2). Compared to cyanide, fluoride




Figure 2. Relative fluorescence emission of **1**·Co (5 μM, CH₃CN) in the presence of various anions (10 μM). I_0 = emission of **1**·Co. I_{CN^-} = fluorescence emission intensity of **1**·Co in the presence of 2 equiv of CN[−]. All the anions were used as tetrabutylammonium salts.

showed a little fluorescence increase even though fluoride is a much more basic anion.¹¹ The nucleophilic azide anion also did not show any significant fluorescence change. These

(4) Bhattacharya, R.; Flora, S. J. S. *Handbook of Toxicology of Chemical Warfare Agents*; Gupta, R. C., Ed.; Academic Press: Boston, 2009; pp 255–270.

(5) (a) Zelder, F. H. *Inorg. Chem.* **2008**, *47*, 1264–1266. (b) Männel-Croisé, C.; Zelder, F. *Inorg. Chem.* **2009**, *48*, 1272–1274. (c) Chow, C.-F.; Lam, M. H. W.; Wong, W.-Y. *Inorg. Chem.* **2004**, *43*, 8387–8393. (d) Kim, Y. H.; Hong, J.-I. *Chem. Commun.* **2002**, 512. (e) Chung, S.-Y.; Nam, S.-W.; Lim, J.; Park, S.; Yoon, J. *Chem. Commun.* **2009**, 2866–2868. (f) Anzenbacher, P., Jr.; Tyson, D. S.; Jursíková, K.; Castellano, F. N. *J. Am. Chem. Soc.* **2002**, *124*, 6232.

(6) (a) Chung, Y.; Lee, H.; Ahn, K. H. *J. Org. Chem.* **2006**, *71*, 9470. (b) Chung, Y. M.; Raman, B.; Kim, D.-S.; Ahn, K. H. *Chem. Commun.* **2006**, 186. (c) Yang, Y.-K.; Tae, J. *Org. Lett.* **2006**, *8*, 5721. (d) Chen, C.-L.; Chen, Y.-H.; Chen, C.-Y.; Sun, S.-S. *Org. Lett.* **2006**, *8*, 5053. (e) Tomasulo, M.; Sortino, S.; White, A. J. P.; Raymo, F. M. *J. Org. Chem.* **2006**, *71*, 744. (f) Lee, K.-S.; Lee, J. T.; Hong, J.-I.; Kim, H.-J. *Chem. Lett.* **2007**, *36*, 816–817. (g) Lee, K.-S.; Kim, H.-J.; Kim, G.-H.; Shin, I.; Hong, J.-I. *Org. Lett.* **2008**, *10*, 49.

results suggest that **1**•Co has a high selectivity for cyanide over other anions.

To address the origin of the fluorescence enhancement of **1**•Co by the coordination of cyanide anions, the HOMO and LUMO energy levels of the cobalt–salen complex in the absence and in the presence of cyanide anions were measured through cyclic voltammetric and UV–vis spectroscopic measurements by using the method of Nagano and co-workers.¹² The energy diagrams indicate that a donor-excited PeT could occur in the absence of cyanide anions, and therefore an electron transfer takes place from the excited coumarin fluorophore to the LUMO of the cobalt–salen complex. The PeT is, however, prohibited in the presence of cyanide anions by the raised LUMO level of the cobalt–salen complex, which can enhance the fluorescence of **1**•Co upon the addition of cyanide (Figure 3). The fluorescence enhancement possibly comes




Figure 3. Energy diagrams (in eV) of coumarin and Co(II)–salen without and with cyanide anion, which was measured from cyclic voltammetry and UV–vis spectroscopy.

from the strong coordination ability of the cyanide anion to **1**•Co.⁵ This strong coordination of cyanide will alter the energy levels in the cobalt(II)–salen complex, especially raising the LUMO level of the cobalt(II)–salen

moiety enough to block the donor-excited PeT and enhance the fluorescence of **1**•Co. However, weak or noncoordinating anions do not complex with the cobalt ion and thus cannot block the donor-excited PeT pathway in **1**•Co. Therefore, the selectivity of **1**•Co plausibly comes from the stronger coordination ability of the cyanide anion over other anions.

As expected, a Job’s plot for binding between **1**•Co and cyanide has shown a 1:2 binding stoichiometry (Supporting Information).¹³ To calculate the binding constants for the 1:2 binding system from the spectral data without assumptions,¹⁴ we have introduced a simple regression program using C language, which is generally applicable to any “one-to-two” binding system (Supporting Information).

Using the program, K_1 and K_2 for cyanide binding to **1**•Co were determined to be $K_1 \geq 10^7 \text{ M}^{-1}$ and $K_2 = 4.0 \times 10^5 \text{ M}^{-1}$ by fluorescence titration in acetonitrile.¹⁵ It is remarkable that the simple cobalt(II)–salen complex shows such a tight binding toward cyanide anions with micromolar dissociation constants. The calculation program shows satisfactory results. The fluorescence intensity constants for HG and HG₂ were calculated to be 166 and 235, respectively, which are very similar to the observed intensities (Figure 4).




Figure 4. Fluorescence titration curve by the addition of cyanide (0–21.9 μM) to **1**•Co (5 μM) in acetonitrile. Calculated (line) and observed data (dot).

In summary, we have developed a Co(II)–salen complex (**1**•Co) which exhibited selective and tight binding to a cyanide anion. **1**•Co displayed a significant fluorescence enhancement upon the addition of cyanide owing to the interruption of photoinduced electron transfer from the coumarin fluorophore to the cobalt(II)–salen moiety. We also developed a general regression method to estimate

(7) Stryer, L. *Biochemistry*, 4th ed.; W. H. Freeman and Company: New York, 1998; pp 60–61 and pp 364–369.

(8) For S100 protein and calcium ion: (a) Marenholz, I.; Heizmann, C. W.; Fritz, G. *Biochem. Biophys. Res. Commun.* **2004**, *322*, 1111–1122. For cAMP and the regulatory subunit of protein kinase A: (b) Weber, I. T.; Takio, K.; Titani, K.; Steitz, T. A. *Proc. Natl. Acad. Sci. U.S.A.* **1982**, *79*, 7679–7683. For transferrin and iron ion: (c) Emerit, J.; Beaumont, C.; Trivin, F. *Biomed. Pharmacother.* **2001**, *55*, 333–339.

(9) (a) Kim, H.-J.; Kim, W.; Lough, A. J.; Kim, B. M.; Chin, J. *J. Am. Chem. Soc.* **2005**, *127*, 16776. (b) Kim, H.-J.; Kim, H.; Alhakimi, G.; Jeong, E. J.; Thavarajah, N.; Studnicki, L.; Koprianiuk, A.; Lough, A. J.; Suh, J.; Chin, J. *J. Am. Chem. Soc.* **2005**, *127*, 16370. (c) Kim, H.; Yen, C.; Preston, P.; Chin, J. *Org. Lett.* **2006**, *8*, 5239. (d) Kim, H.-J.; Chin, J.; Lough, A. J. *Acta Crystallogr.* **2007**, *E63*, o3901. (e) Kim, H.; Yen, C.; Preston, P.; Chin, J. *Org. Lett.* **2006**, *8*, 5239. (f) Kim, H.; Nguyen, Y.; Yen, C. P.-H.; Chagal, L.; Lough, A. J.; Kim, B. M.; Chin, J. *J. Am. Chem. Soc.* **2008**, *130*, 12184.

(10) See the Supporting Information.

(11) Bordwell, F. G. *Acc. Chem. Res.* **1988**, *21*, 456.

(12) (a) Tanaka, K.; Miuram, T.; Umezawa, N.; Urano, Y.; Kikuchi, K.; Higuchi, T.; Nagano, T. *J. Am. Chem. Soc.* **2001**, *123*, 2530–2536. (b) Ueno, T.; Urano, Y.; Setsukinai, K.; Takakusa, H.; Kojima, H.; Kikuchi, K.; Ohkubo, K.; Fukuzumi, S.; Nagano, T. *J. Am. Chem. Soc.* **2004**, *126*, 14079–14085.

(13) Connors, K. A. *Binding Constants*, 1st ed.; John Wiley & Sons: New York, 1987; pp 24–28.

(14) (a) Kavallieratos, K.; Bertao, C. M.; Crabtree, R. H. *J. Org. Chem.* **1999**, *64*, 1675. (b) Hynes, M. J. *J. Chem. Soc., Dalton Trans.* **1993**, 311. (c) Nishizawa, S.; Kaneda, H.; Kato, Y.; Teramae, N. *J. Am. Chem. Soc.* **1999**, *121*, 9463.

(15) Using the program, K_1 was calculated over 10^7 M^{-1} in acetonitrile, which is above the limit of detection. Detailed programming for the calculation of K_1 and K_2 is described in the Supporting Information.

the binding constants in the 1:2 binding system and found the dissociation constants of **1•Co** to be in the range of micromolar concentrations for cyanide anions.

Acknowledgment. This work was supported by the KRF Grant (Grant No. 2008-314-C00206) and the NRF grant funded by the MEST (Grant No. 2009-0080734). ARJ and ISS thank the Ministry of Education for the BK 21 fellowship.

Supporting Information Available: Synthetic procedure and selected spectral data for compounds **1** and **1•Co**, fluorescence spectra, Job's plots, electrochemical data, and a regression program for the determination of binding constants. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL902852G